Conger myriaster, a new second intermediate host of *Heterophyopsis continua* (Digenea: Heterophyidae)

Ki-Hong KIM^{*1)}, Eun-Seok CHOI¹⁾ and Han-Jong RIM²⁾

Department of Fish Pathology¹⁾, College of Fisheries Sciences, Pukyong University, Pusan 608-737,
Department of Parasitology²⁾ and Institute for Tropical Endemic Diseases, College of Medicine,
Korea University, Seoul 136-705, Korea

Abstract: Six metacercariae were found from the gill filaments of *Conger myriaster* purchased at Mokpo-shi in Korea on 7 September, 1996. Based on the morphology of the excysted specimen, we identified them as metacercariae of *Heterophyopsis continua*. *C. myriaster* is a new intermediate host of *H. continua* in the literature.

Key words: *Heterophyopsis continua*, metacercaria, the second intermediate host, *Conger myriaster*

Heterophyopsis continua is a species of family Heterophyidae and is parasitic in fish eating birds and mammals. Three human cases of natural infection by *H. continua* were recorded in Korea (Seo *et al.*, 1984; Hong *et al.*, 1996).

Brackish water or marine fishes — *Lateolabrax japonicus*, *Mugil cephalus*, *Acanthogobius flavimanus*, *Plecoglossus altivelis* and *Clupanodon punctatus* — were recorded as the second intermediate hosts of *H. continua* in Korea (Chun, 1960; Seo *et al.*, 1984; Cho and Kim, 1985; Sohn *et al.*, 1994).

On 7 September, 1996, we purchased several species of marine fish in a fisheries market in Mokpo city in order to investigate helminth parasites. We found 6 metacercariae in the gill filaments of two conger eels (*Conger myriaster*). Among those metacercariae, one excysted metacercaria was fixed in 70% hot ethylalcohol. The fixed specimen was stained with Semichon's acetocarmine.

Morphology of the stained specimen is as follow; Body elongated leaf-like, dorsoventrally

flattened, 1.260 mm long and 0.230 mm wide (Fig.1). Tegument beset with small spines which is sparse posteriorly. Oral sucker subterminal, 0.065 x 0.078 mm. Ventral sucker 0.120 x 0.113 mm, lying at the slightly anterior to the half of body. Prepharynx very long, 0.176 mm in length. Pharynx elliptical, 0.075 x 0.050 mm. Oesophagus very short. Intestinal caeca extended to posterior end of body. Genital sucker larger than oral sucker but smaller than ventral sucker, 0.078 x 0.100 mm, situated slightly posterosinistral to ventral sucker. Rodlets on the genital sucker were all the same size, approximately 100 in number, and arranged in a single circle. Ovary elliptical, 0.028 x 0.043 mm. Two testes smooth, globular and a little obliquely tandem. Anterior testis 0.055 x 0.053 mm and posterior testis 0.060 x 0.050 mm. Excretory vesicle Y-shaped, extended to the posterior margin of ovary.

The characteristics of this specimen was coincide with those of *Heterophyopsis continua*. There was no report that *C. myriaster* acts as the second intermediate host of *H. continua*. We, therefore, report here *C. myriaster* as a fish intermediate host of *H.*

• Received Nov. 9 1996, accepted Nov. 29 1996.

*Corresponding author

Fig. 1. Diagram of excysted metacercaria of *Heterophyopsis continua* collected from the gill filaments of *Conger myriaster* (ventral view, scale bar = 0.3 mm)

continua for the first time in the world.

Generally, the infection environment of *H. continua* to fish was considered via brackish water near river mouth, but *C. myriaster* and *C. punctatus* are not brackish water fish but coastal water fish. Therefore it is possible that the first intermediate hosts of *H. continua* may be the euryhalne organisms which can survive both in brackish water and sea water.

REFERENCES

- Cho SY, Kim SI (1985) *Plecoglossus altivelis* as a new fish intermediate host of *Heterophyopsis continua*. *Korean J Parasitol* **23**(1): 173-174.
- Chun SK (1960) A study on some trematodes whose intermediate hosts are brackish water fish (1) The life history of *Heterophyes continus* the intermediate host of *Laterolabrax japonicus*. *Bulletin of Pusan Fisheries College* **3**(1&2): 40-44 (in Korean).
- Hong SJ, Chung CK, Lee DH, Woo HC (1996) One human case of natural infection by *Heterophyopsis continua* and three other species of intestinal trematodes. *Korean J Parasitol* **34**(1): 87-90.
- Seo BS, Lee SH, Chai JY, Hong SJ (1984) Studies on intestinal trematodes in Korea XIII. Two cases of natural human infection by *Heterophyopsis continua* and the status of metacercarial infection in brackish water fishes. *Korean J Parasitol* **22**(1): 51-60.
- Sohn WM, Han GG, Kho WG, Chai JY, Lee SH (1994) Infection status with the metacercariae of heterophyid flukes in the brackishwater fish from Haenam-gun, Chollanam-do, Korea. *Korean J Parasitol* **32**(3): 163-169 (in Korean).

=초록=

긴이형흡충(*Heterophyopsis continua*)의
새로운 중간숙주인 붕장어(*Conger myriaster*)

김기홍¹⁾, 최은석¹⁾, 임한중²⁾

부경대학교 수산과학대학 어병학과¹⁾ 및 고려대학교 의과대학 기생충학교실²⁾

1996년 9월 7일 전남 목포항에서 구입한 붕장어의 아가미 새엽에서 6개체의 피낭유충이 검출되었다. 이중 탈낭된 한 개체의 형태를 관찰한 바 긴이형흡충의 피낭유충으로 동정되었으며, 붕장어가 긴이형흡충의 제2중간숙주 역할을 한다는 것을 처음으로 보고한다.

[기생충학잡지 34(4): 283-285, 1996년 12월]