Cloning of a pore-forming subunit of ATP-sensitive potassium channel from *Clonorchis sinensis*

Seung-Young HWANG¹⁾, Hye-Jin HAN¹⁾, So-Hee KIM¹⁾, Sae-Gwang PARK²⁾, Dae-Hyun SEO^{2),4)},
Nari KIM^{3),4)}, Jin HAN^{3),4)}, Joon-Yong CHUNG^{1),4)}* and Weon-Gyu KHO¹⁾

¹⁾Department of Parasitology and Institute of Malariology, ²⁾Department of Microbiology, ³⁾Department of Physiology and Biophysics, ⁴⁾Molecular Cell Physiology Research Group, Inje University College of Medicine, Busan 614-735, Korea

Abstract: A complete cDNA sequence encoding a pore-forming subunit (Kir6.2) of ATP-sensitive potassium channel in the adult worm, *Clonorchis sinensis*, termed CsKir6.2, was isolated from an adult cDNA library. The cDNA contained a single open-reading frame of 333 amino acids, which has a structural motif (a GFG-motif) of the putative pore-forming loop of the Kir6.2. Peculiarly, the CsKir6.2 shows a lack-sequence structure, which deleted 57 amino acids were deleted from its N-terminus. The predicted amino acid sequence revealed a highly conserved sequence as other known other Kir6.2 subunits. The mRNA was weakly expressed in the adult worm.

Key words: *Clonorchis sinensis*, potassium channels, cloning

Adenosine 5'-triphosphate-sensitive potassium channels (K_{ATP} channels) are thought to regulate various cellular functions such as secretion as well as muscular and neural excitability by linking the cell's metabolic state to its membrane potential (Noma, 1983; Bernardi et al., 1993). K_{ATP} channels were originally discovered in the heart (Noma, 1983) and found later in various tissues, including pancreatic-cells (Rorsman et al., 1985), smooth and skeletal muscles (Standen et al., 1989), the brain (Ashford et al., 1988), pituitary glands (Bernardi et al., 1988), and the kidney (Hunter et al., 1988). It has been discovered that the K_{ATP} channels are comprised of two kinds of subunits: sulphonylurea receptors (SUR) which belong to the ATP-binding cassette family and

inwardly-rectifying K^+ channel (Kir) subunits which forms the potassium ion-selective pore. The channels are proposed to form a complex of SUR1 and Kir6.2 in pancreatic cells (Sakura et al., 1995; Inagaki et al., 1996), and SUR2A and Kir6.2 in cardiac and skeletal muscles (Inagaki et al., 1996). In particular, several Kir6.2 genes have been identified in human, mouse, rat and rabbit tissues, showing high mRNA levels in the heart, skeletal muscle, bladder and gut. Therefore, the K_{ATP} channel of the helminth worm is thought to be important in the cellular communication and metabolic pathway. However, the information on the study of the K_{ATP} channel for the helminth worm has been scarce until now.

To add useful information for ion channel of the helminth worm, the gene encoding a pore-forming subunit of the liver fluke in the adult worm *Clonorchis sinensis* (CsKir6.2) was isolated and characterized. In addition, the gene expression level of a CsKir6.2 in the adult worm was investigated using a Northern blot

• Received 21 April 2003, accepted after revision 27 May 2003.

• This work was supported by the 2002 Inje University research grant.

*Corresponding author (e-mail: paracjyo@inje.ac.kr)

analysis.

The total RNA was isolated from the adult of *C. sinensis* by guanidium isothiocyanate lysis and CsCl centrifugation at $32,000 \times g$ for 16 hr at 20°C . A cDNA library was constructed using SMART cDNA library

synthesis kit (Clontech., Palo Alto, CA, USA). Briefly, the first-strand cDNAs were synthesized using PowerScript reverse transcriptase, SMART IV oligonucleotide and a modified oligo (dT) primer (CDS III/3' PCR primer). The second-strand cDNAs

Fig. 1. Nucleotide and deduced amino acid sequences of a pore-forming subunit (Kir6.2) of K_{ATP} channel from *Clonorchis sinensis*. The single underline shows the region corresponding to the putative transmembrane, and the dotted underline shows pore domains. An asterisk indicates a stop codon. The bold letter shows a GFG-motif (the putative pore-forming loop). The position of primers for cDNA PCR (Kir6.2F and Kir6.2R) and gDNA PCR (CsKir6.2F and CsKir6.2R) indicated above the nucleotide sequence.

were synthesized using 50 × Advantage 2 polymerase Mix, 5' PCR primer, and a CDS/3' PCR primer. The resulting cDNA was digested with *Sfi* I, and fractionated using Chroma Spin-400. The cDNAs larger than 0.5 kb were pooled (data not shown). After digestion with *Sfi* I and size fractionation, the cDNA was ligated into the *Sfi* I-digested λ Triplex2 vector using T4 DNA ligase. The ligated cDNAs were packaged in Gigapack III gold packaging extract (Stratagene, La Jolla, CA, USA). To obtain a partial fragment of a specific probe, 1 gm of total RNA from the adult worm *C. sinensis* was reverse-transcribed using an oligo d(T)₁₅ primer and Moloney Murine Leukemia Virus (M-MuLV) reverse transcriptase (Life Technologies, Gaithersburg, MD, USA). Two degenerate oligonucleotides (Kir6.2F, 5-TGACCTGGCSCCCRGCGA-3 and Kir6.2R, 5-CRTCKGC CAGGTAGGAGGT-3 S, G+C; R, A+G; K, G+T) were designed and synthesized according to the highly conserved regions of various human and rabbit ATP-sensitive inwardly-rectifying potassium channels, termed Kir6.2. The resulting cDNA was subjected to PCR amplification in a 50 μ l reaction mixture containing 5 μ l of RT-PCR product, 5 μ l of 10 × *Taq* polymerase buffer, 4 μ l of 2.5 mM dNTPs, 25 pM of each primers and 5 units of *Taq* polymerase. The reaction was subjected to 35 cycles of denaturation at 95°C for 30 sec, annealing at 55°C for 1 min, and extension at 72°C for 30 sec, and one at 72°C for 5 min in a thermal cycler (GeneAmp PCR system 9700, Perkin-Elmer). PCR with these primers and the first strand cDNA from the total RNA amplified an approximately 620 bp cDNA fragment, which then was sequenced by direct sequencing. The PCR fragment was showed the significant sequence homology with Kir 6.2 family (data not shown). An adult worm *C. sinensis* cDNA library of approximately 100,000 independent plaques was screened by plaque hybridization employing 620 bp partial fragment, which was amplified by RT-PCR. Hybridization to the ³²P-labeled probe was done overnight at 42°C. The membrane was washed with high stringency and exposed to Kodak X-AR film. Positive clones were isolated and the lambda DNA was purified by use of a Qiagen Lambda kit (Qiagen, Valencia, CA). The cDNA insert was cloned into a pGEM-T easy vector

(Promega) by PCR, using λ Triplex Sequencing primer set (Clontech) and advanced *Taq* polymerase with proofreading capacity (Clontech). The recombinant plasmid in the bacterial culture was purified with a Qiagen plasmid minikit (Qiagen). The nucleotide sequence was determined by the Dideoxy Chain Termination method using a Sequenase kit (ABI Prism Dye Terminator Cycle Sequencing Core Kit, Perkin Elmer) and an automated DNA sequencer (Applied Biosystems model 377A; Perkin Elmer). The determined nucleotide sequence and deduced amino acid sequence were analyzed with Basic Local Alignment Search Tool (BLAST) in National Center for Biotechnology Information (NCBI) and ExPASy Molecular Biology Server (<http://www.expasy.ch>).

The one clone containing the largest insert (1,189 bp in length) revealed a 999 bp complete open reading frame (ORF) coding for a protein of 333 amino acids. The 3'-untranslated region was contained poly (A)⁺ tail (Fig. 1). This sequence has been deposited in the GenBank database with the accession number AY277926. A hydrophobicity plot of the Kir6.2 revealed two putative transmembrane domains as found in other Kir channels. In addition, a GFG-motif was found in the putative pore-forming loop. The GFG-motif was showed in only both Kir6.1 and Kir6.2 (Inagaki et al., 1996). Although Kir6.2 shows a high degree of homology with Kir6.1, the intracellular N-terminal and C-terminal regions and an extracellular region between the first transmembrane region and the pore region (H5) do not. Therefore, the nucleotide sequence of CsKir6.2 is more similar to other known Kir6.2 than to other ion channel subunits. However, the CsKir6.2 has more unusual structural features than do other Kir6.2 genes (Fig. 2). The CsKir6.2 is smaller than Kir6.2 subunits of other species, containing a lacked 57 amino acids sequence in the N-terminal region.

In the case of ruling out a lacked sequence region, a comparison of the amino acid sequence for CsKir6.2 with the Kir6.2 subunits of other species in the database showed a high identity to the proteins of human, mouse, rabbit, guinea pig and rat (Fig. 2). There was a difference of 13, 14, 6, 20, and 13 amino acids between the sequences of *C. sinensis* and the guinea pig, human, mouse, rat, and rabbit,

respectively. A Northern blot was used to investigate the expression of Kir6.2 in *C. sinensis*. A single RNA transcript of approximately 1 kb in size were weakly expressed (Fig. 3A). In order to reveal the cause of a lacked-sequence structure in the Cskir6.2, it was

amplified against genomic DNA using CsKir6.2F (5'-ATGCTGTCCCGAAAGGGCAT-3') and CsKir6.2R (5'-TCAGGACAAGGAATCTGGAG-3') primers. Samples were denatured at 95°C for 2 min, followed by 35 cycles of 94°C for 30 sec, 58°C for 30 sec and

<i>C. sinensis</i>	MLSRKGIIPEEYVLTRLAEDPAEP-----	24
Rabbit	MLSRKGIIPEEYVLTRLAEDPAEPYRARRERRARFVSKKGCNVAHKNIREQGRFLQDVF	60
Rat	MLSRKGIIPEEYVLTRLAEDPTEPRYRTRERRARFVSKKGCNVAHKNIREQGRFLQDVF	60
Mouse	MLSRKGIIPEEYVLTRLAEDPAEPYRTRERRARFVSKKGCNVAHKNIREQGRFLQDVF	60
Human	MLSRKGIIPEEYVLTRLAEDPAKPRYRARRRRARFVSKKGCNVAHKNIREQGRFLQDVF	60
Guineapig	MLSRKGIIPEEYVLTRLAEDPTEPRYRARRERRARFVSKKGCNVAHKNIREQGRFLQDVF	60
	***** *	
<i>C. sinensis</i>	-----SWLLFAMVWWLIAFAHGD LAPGEGA AVPCVTSIHSLSA	63
Rabbit	TTLVDLKWHTLLIFTMSFLCSWLLFAMVWWLIAFAHGD LAPGEGA AVPCVTSIHSFS	120
Rat	TTLVDLKWPHHTLLIFTMSFLCSWLLFAMVWWLIAFAHGD LAPGEGTNVPCVTSIHSFS	120
Mouse	TTLVDLKWPHHTLLIFTMSFLCSWLLFAMVWWLIAFAHGD LAPGEGTNVPCVTSIHSFS	120
Human	TTLVDLKWPHHTLLIFTMSFLCSWLLFAMAWWLIAFAHGD LAPSEGTAEPVTSIHSFS	120
Guineapig	TTLVDLKWPHHTLLIFTMSFLCSWLLFAMVWWLIAFAHGD LAPGEGTTPCVTSIHSFS	120
	***** ** *	
<i>C. sinensis</i>	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTAQAHRRAET	123
Rabbit	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTAQAHRRAET	180
Rat	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTAQAHRRAET	180
Mouse	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTAQAHRRAET	180
Human	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTAQAHRRAET	180
Guineapig	FLFSIEVQVTIGFGGRMVTEECPLAILILIVQNIVGLMINAIMLGCI FMKTSQAHRRAET	180
	***** *	
<i>C. sinensis</i>	LIFS KHAVIALRQDRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	183
Rabbit	LIFS KHAVIALRQGRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	240
Rat	LIFS KHAVITLRHGRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	240
Mouse	LIFS KHAVITLRHGRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	240
Human	LIFS KHAVIALRHGRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	240
Guineapig	LIFS KHAVIALRHGRLCFMLRVGDLRKSMIISATIHMQVVRKTTSPGEV VPLHQVDIPM	240
	***** ** *	
<i>C. sinensis</i>	ENGVGGSIFLVAPLI IHHVIDTNSPLYDLAPSDLHHHQDLEI IVILEGVVETTGITTTQA	243
Rabbit	ENGVGGSIFLVAPLI IHHVIDANSPLYDLAPSDLHHHQDLEI IVILEGVVETTGITTTQA	300
Rat	ENGVGGSIFLVAPLI IYHVIDSNSPLYDLAPSDLHHHQDLEI IVILEGVVETTGITTTQA	300
Mouse	ENGVGGSIFLVAPLI IYHVIDSNSPLYDLAPSDLHHHQDLEI IVILEGVVETTGITTTQA	300
Human	ENGVGGSIFLVAPLI IYHVIDANSPLYDLAPSDLHHHQDLEI IVILEGVVETTGITTTQA	300
Guineapig	ENGVGGSIFLVAPLI IYHVIDANSPLYDLGPSDLHHHQDLEI IVILEGVVETTGITTTQA	300
	***** *	
<i>C. sinensis</i>	RTSYLADEILWGQRFVPIVAEEDGRYSVDYSKFGNAV KVP TPLCTARQLDEDRSLLDALT	303
Rabbit	RTSYLADEILWGQRFVPIVAEEDGRYSVDYSKFGNTVKVP TPLCTARQLDEDRSLLDALT	360
Rat	RTSYLADEILWGQRFVPIVAEEDGRYSVDYSKFGNTVKVP TPLCTARQLDEDRSLLDALT	360
Mouse	RTSYLADEILWGQRFVPIVAEEDGRYSVDYSKFGNTIKVP TPLCTARQLDEDRSLLDALT	360
Human	RTSYLADEILWGQRFVPIVAEEDGRYSVDYSKFGNTVKVP TPLCTARQLDEDRSLLEALT	360
Guineapig	RTSYLADEILWGHRFVPIVAEEDGRYSVDYSKFGNTIKVP TPLCTAHQLDEDRSLLDALT	360
	***** ** *	
<i>C. sinensis</i>	LTSARGSLRKRSVAVAKAKPKFSISPDSLS	333 % identity
Rabbit	LTSARGPLRKRSVPVAKAKPKFSISPDSLS	390 98.2
Rat	LASSRGPLRKRSVAVAKAKPKFSISPDSLS	390 96.1
Mouse	LASSRGPLRKRSVAVAKAKPKFSISPDSLS	390 95.8
Human	LASARGPLRKRSVPMKAKPKFSISPDSLS	390 95.2
Guineapig	LASTRGPLRKRSVPVAKAKPRFSISPDSLS	390 94.0
	* * * * *	

Fig. 2. Comparison of deduced amino acid sequences of *Clonorchis sinensis* Kir6.2 (CsKir6.2), rabbit (*Oryctolagus cuniculus*, the Genbank No. D45025), rat (*Rattus norvegicus*, the Genbank No. BAA96239), mouse (*Mus musculus*, the Genbank No. Q61743), human (*Homo sapiens*, the Genbank accession number Q14654), and guinea pig (*Cavia porcellus*, the Genbank No. Q9JHJ9). Gaps indicated by dashes (- -) are introduced to optimize the alignment. An asterisk indicates an amino acid that all peptides show identical one to the query sequence.

Fig. 3. Northern blot analysis and genomic DNA amplification of *CsKir6.2*. (A) The total RNA from adult *Clonorchis sinensis* was separated on a 1.2% formaldehyde agarose gel and transferred to a nylon membrane. The labeling of the probe and detection of hybridization signal were carried out with a ^{32}P -labelled system. (B) The *CsKir6.2* was amplified against the genomic DNA and cDNA of adult *C. sinensis*, then separated on a 1% agarose gel. Lane M, $\gamma\text{DNA}/\text{EcoR I} + \text{Hind III}$ marker (MBI fermentas); lane 1, PCR-amplified genomic DNA; lane 2, PCR amplified cDNA.

72.5°C for 1 min. A 5 min final extension was performed at 72.5°C. PCR products were analyzed on 1% agarose gel. The nucleotide sequence was determined by dideoxynucleotide chain termination method using a sequenase kit (ABI PRISM Dye Terminator Cycle Sequencing Core Kit) and an automated DNA sequencer. Amplicon of PCR was identical to the cDNA in size (Fig. 3B) and nucleotide sequence (data not shown). Therefore, the lacked-sequence structure of *CsKir6.2* is thought to be a peculiar characteristic of *C. sinensis* and is not isoform.

In this study, a pore-forming subunit of the K_{ATP} channel (Kir6.2 gene) of *C. sinensis* was cloned. This is the first report on the cloning of the Kir6.2 gene from trematodes. These results demonstrate that Kir6.2 encodes an inwardly-rectifying KATP channel protein

which is expressed in adult *C. sinensis*. Generally, the K_{ATP} channel consists of a pore forming subunit (Kir6.2) and a sulphonylurea receptor (SUR). Both subunits are required to form a functional K_{ATP} channel. Therefore, further study is needed to address the exact functions of K_{ATP} channel by co-expressing Kir6.2 and SUR after the SUR gene is cloned in *C. sinensis*.

REFERENCES

- Ashford ML, Sturgess NC, Trout NJ, Gardner NJ, Hales CN (1988) Adenosine-5'-triphosphate-sensitive ion channels in neonatal rat cultured central neurons. *Eur J Physiol* **412**: 297-304.
- Bernardi H, Fosset M, Lazdunski M (1988) Characterization, purification, and affinity labeling of the brain [^3H]glibenclamide-binding protein, a putative neuronal ATP-regulated K^+ channel. *Proc Nat Acad Sci USA* **85**: 9816-9820.
- Bernardi H, De-Weille JR, Epelbaum J, et al. (1993) ATP-modulated K^+ channels sensitive to antidiabetic sulfonylureas are present in adenohypophysis and are involved in growth hormone release. *Proc Nat Acad Sci USA* **90**: 1340-1344.
- Inagaki N, Gonoi T, Clement JP, et al. (1996) A family of sulfonylurea receptors determines the pharmacological properties of ATP-sensitive K^+ channels. *Neuron* **16**: 1011-1017.
- Noma A (1983) ATP-regulated K^+ channels in cardiac muscle. *Nature* **305**: 147-148.
- Rorsman P, Trube G (1985) Glucose dependent K^+ channels in pancreatic-cells are regulated by intracellular ATP. *Eur J Physiol* **405**: 305-309.
- Sakura H, Ammala C, Smith PA, Gribble FM, Ascroft FM (1995) Cloning and functional expression of the cDNA encoding a novel ATP-sensitive potassium channel subunit expressed in pancreatic-cells, brain, heart and skeletal muscle. *FEBS Lett* **377**: 338-344.
- Standen NB, Quayle JM, Davies NW, Brayden JE, Huang Y, Nelson MT (1989) Hyperpolarizing vasodilators activate ATP-sensitive K^+ channels in arterial smooth muscle. *Science* **245**: 177-180.